

Parthenium weed

Parthenium hysterophorus



Parthenium costs the beef industry a total of \$16.5 million per year and cropping industries several million dollars per year.

Declaration details

In Queensland, Parthenium is a Class 2 declared plant.

Under the *Land Protection (Pest and Stock Route Management) Act 2002*, Class 2 declaration requires landholders to control pests on the land and waters under their control. A local government may serve a notice upon a landholder requiring control of declared pests.



Queensland Government

Description and general information

Size

Parthenium weed is an annual herb with a deep tap root and an erect stem that becomes woody with age. As it matures, the plant develops many branches in its top half and may eventually reach a height of two metres.

Leaves

Its leaves are pale green, deeply lobed and covered with fine soft hairs.

Flowers

Small creamy white flowers occur on the tips of the numerous stems. Each flower contains four to five black seeds that are wedge-shaped, two millimetres long with two thin, white scales.

Lifecycle

Parthenium weed normally germinates in spring and early summer, produces flowers and seed throughout its life and dies around late autumn. However, with suitable conditions (rain, available moisture, mild temperatures), parthenium weed can grow and produce flowers at any time of the year. In summer, plants can flower and set seed within four weeks of germination, particularly if stressed.

Potential damage

Parthenium weed is a vigorous species that colonises weak pastures with sparse ground cover. It will readily colonise disturbed, bare areas along roadsides and heavily stocked areas around yards and watering points. Parthenium weed can also colonise brigalow, gidgee and softwood scrub soils. Its presence reduces the reliability of improved pasture establishment and reduces pasture production potential.

Parthenium weed is also a health problem as contact with the plant or the pollen can cause serious allergic reactions such as dermatitis and hay fever.

Habitat and distribution

Parthenium weed is capable of growing in most soil types but becomes most dominant in alkaline, clay loam soils.

The plant is well established in Central Queensland and present in isolated infestations west to Longreach and in northern and southern Queensland.

Infestations have also been found in northern and central parts of New South Wales and it is capable of growing in most states of Australia.

Control

Prevention and weed seed spread

As with most weeds, prevention is much cheaper and easier than cure. Pastures maintained in good condition, with high levels of grass crown cover, will limit parthenium

weed colonisation. Drought, and the subsequent reduced pasture cover, creates the ideal window of opportunity for parthenium weed colonisation when good conditions return.

Parthenium seeds can spread via water, vehicles, machinery, stock, feral and native animals and in feed and seed. Drought conditions aid the spread of seed with increased movements of stock fodder and transports.

Vehicles and implements passing through parthenium weed infested areas should be washed down with water. Wash down facilities are located in Alpha, Biloela, Charters Towers, Emerald, Gracemere, Injune, Monto, Moura, Rolleston, Springsure and Taroom. Particular care should be taken with earthmoving machinery and harvesting equipment. The wash down procedure should be confined to one area, so that plants that establish from dislodged seed can be destroyed before they set seed.

Extreme caution should be taken when moving cattle from infested to clean areas. Avoid movement during wet periods as cattle readily transport seed in muddy soil. On arrival, cattle should be held in yards or small paddocks until seed has dropped from their coats and tails prior to their release into large paddocks. Infestations around yards can be easily spotted and controlled whereas infestations can develop unnoticed in large paddocks.

Particular care should be taken when purchasing seed, hay and other fodder materials. Always keep a close watch on areas where hay has been fed out for the emergence of parthenium or other weeds.

Property hygiene is important. Owners of clean properties should ensure that visitors from infested areas do not drive through their properties. If your property has parthenium weed on it, ensure that it is not spread beyond the boundary or further within the property.

Pasture management

Grazing management is the most useful method of controlling large-scale parthenium weed infestations. Maintain pastures in good condition with high levels of ground and grass crown cover. This may require rehabilitation of poor pastures, followed by a sound grazing maintenance program.

Sown pasture establishment—Poor establishment of sown pastures can allow parthenium weed colonisation. pasture agronomist Aerial seeding prior to scrub pulling is normally beneficial.

Overgrazing—High grazing pressure caused by drought or high stock numbers decreases the vigour and competitiveness of pastures and allows the entry and spread of parthenium weed. Maintenance of correct stock numbers is most important in controlling parthenium weed. pasture agronomist

Pastures spelling—In situations of serious infestation, pasture spelling is essential for rehabilitation. Total spelling is much more effective than simply reducing the

stocking rate. However, overgrazing of the remainder of the property must be avoided.

The most appropriate time for pasture spelling is the spring–summer growing period, with the first 6–8 weeks being particularly important. If the condition of perennial grasses (native or sown) is low, spelling for the entire growing season may be required or introduced grasses may need to be re-sown. Herbicide treatment can hasten the rehabilitation process by removing a generation of parthenium seedlings and allowing grass seedlings to establish without competition. In the presence of parthenium weed, grass establishment is poor.

Grazing during winter should not increase the parthenium weed risk. Most tropical grasses are dormant and can tolerate moderate grazing during this period. However, parthenium weed may germinate and grow at this time.

Fencing—One of the main problems in controlling parthenium weed is the large paddock size and the variability of country within paddocks. The resulting uneven grazing pressures encourage parthenium weed to colonise the heavily grazed country. Ideally, similar land types should be fenced as single units. Fencing can be used to great effect to break up large paddocks, allowing more flexible management such as pasture spelling or herbicide application, options not available previously.

Burning—Burning is not promoted as a control strategy for parthenium weed. However, research suggests that burning for pasture management (e.g. woody weed control) should not result in an increased infestation if the pasture is allowed to recover prior to the resumption of grazing. Stocking of recently burnt areas known or suspected to contain parthenium decreases pasture competition and favours parthenium, ultimately creating a more serious infestation.

Herbicide control

Non-crop areas—Parthenium weed should be sprayed early before it can set seed. A close watch should be kept on treated areas for at least two years.

Small and/or isolated infestations should be treated immediately. Herbicide control will involve a knockdown herbicide to kill plants that are present and a residual herbicide to control future germinations. Repeated spraying may be required even within the one growing season to prevent further seed production.

Extensive infestations will require herbicide treatment in conjunction with pasture management. Timing of spraying is critical so that parthenium weed is removed when plants are small and before seeding has occurred. Grasses should be actively growing and seeding so that they can recolonise the infested area.

Table 1 shows the herbicides registered for parthenium weed control and application rates. Before using any herbicide always read the label carefully. All herbicides must be applied strictly in accordance with the directions on the label.

Cropping areas—Controlling parthenium weed in cropland requires selective herbicide use and/or crop rotations. For further information on parthenium weed control in crops consult your nearest Primary Industries and Fisheries extension agronomist.

Biological control

The combined effects of biological control agents reduced the density and vigour of parthenium weed and increased grass production.

There are currently a number of insect species and two rust pathogens that have been introduced to control parthenium weed—a selection of these are outlined below.

Epiblema strenuana is a moth introduced from Mexico established in all parthenium weed areas. The moth's larvae feed inside the stem, forming galls that stunt the plant's growth, reduce competitiveness and seed production.

Listronotus setosipennis is a stem-boring weevil from Argentina but is of limited success in reducing parthenium weed infestations.

Zygogramma bicolorata is a defoliating beetle from Mexico which is highly effective where present. It emerges in late spring and is active until autumn.

Smicronyx lutulentus (Mexico) lays eggs in the flower buds where the larvae feed on the seed heads.

Conotrachelus albocinereus (stem-galling weevil from Argentina) produces small galls and is still becoming established in Queensland.

Bucculatrix parthenica (leaf mining moth from Mexico) larvae feed on leaves, leaving clear windows in the leaf.

Carmentia ithacae is a stem boring moth from Mexico which is becoming established at favourable sites in the northern Central Highlands.

Puccinia abrupta is a winter rust from Mexico that infects and damages leaves and stems. It is currently established over a wide area from Clermont south. It requires a night temperature of less than 16 degrees and 5–6 hours of leaf wetness (dew). Sporadic outbreaks occur where weather conditions are suitable.

Puccinia melampodii is a summer rust from Mexico that weakens the plant by damaging the leaves over the summer growing season. It is currently established and spreading at a number of sites from north of Charters Towers to Injune in the south.

Manual control

Hand pulling of small areas is not recommended. There is a health hazard from allergic reactions and a danger that mature seeds will drop off and increase the area of infestation.

Further information

Further information is available from your local government office, or from your local primary industries and fisheries biosecurity officer: contact details are available through 13 25 23.

Table 1 Herbicides registered for parthenium weed.

Herbicide	Rate	Situation	Comments
2,4-D amine 500 g/L	0.4 L/100 L	Land—industrial, pastures; rights-of-way	Spot spray
atrazine 500 g/L max 3 kg/ha/yr	3.6–6 L/ha	Fields and fallow	Boom spray
	6 L/ha	Land—industrial, commercial, non-agricultural, roadside, right-of-way	Boom spray
atrazine 900 g/kg max 3 kg/ha/yr	2–3.3 kg/ha	Fields and fallow	Boom spray
	3.3 kg/ha	Land—non-agricultural, commercial, industrial	Boom spray
2,4-D + picloram (Tordon 75-D)	125 ml/100 L	Land—commercial, industrial, pastures, right-of-way	Spot spray
	3 L/ha	Land—commercial, industrial, pastures, right-of-way	Boom spray
2,4-D ester ¹	.025 L/10 L	Land—non-agricultural, pastures	Rosette stage
glyphosate (450 g/L)	0.8–1.2 L/ha	Fields and fallow	Spot spray
metsulfuron methyl	5–7 g/ha	Fields and fallow	Seedlings only
	5 g/100 L	Land—commercial, industrial, pastures, rights-of-way	Spot spray
hexazinone	3.5 L/ha or 7 L/10 L/20 m ²	Land—commercial, industrial, pastures, rights-of-way	Boom spray or spot spray
dicamba (200 g/L) (500 g/L) (700 g/kg)	0.7–2.8 L/ha or 0.1–0.19 L/100L	Grass pastures	Boom spray or spot spray
	0.28–1.1 L/ha or 0.40–0.76 L/100L	Grass pastures	Boom spray or spot spray
	200–800 g/ha or 30–60 g/100 L	Grass pastures	Boom spray or spot spray

¹Use restricted in some areas of Central Queensland

Notes The registered rates are for non-crop uses. Consult label for in-crop recommendations.
For power hand spray or knapsack use, spray plants to the point of runoff.

Fact sheets are available from Queensland Primary Industries and Fisheries service centres and the Queensland Primary Industries and Fisheries Business Information Centre (telephone 13 25 23). Check our website at www.dpi.qld.gov.au to ensure you have the latest version of this fact sheet. The control methods referred to in this pest fact should be used in accordance with the restrictions (federal and state legislation, and local government laws) directly or indirectly related to each control method. These restrictions may prevent the use of one or more of the methods referred to, depending on individual circumstances. While every care is taken to ensure the accuracy of this information, Queensland Primary Industries and Fisheries does not invite reliance upon it, nor accept responsibility for any loss or damage caused by actions based on it.